arXiv:1406.6924v1 [cs.SC] 26 Jun 2014

STRONGLY STABLE IDEALS AND HILBERT POLYNOMIALS

DAVIDE ALBERELLI AND PAOLO LELLA

Abstract. The *StronglyStableIdeals* package for *Macaulay2* provides a method to compute all the saturated strongly stable ideals in a given polynomial ring defining schemes with a fixed Hilbert polynomial. A description of the main method and the auxiliary tools is given.

Introduction

Strongly stable ideals are a key tool in commutative algebra and algebraic geometry. These ideals have nice combinatorial properties that makes them well suited for both theoretical and computational applications. In the case of polynomial rings with coefficients in a field with characteristic zero, the notion of strongly stable ideals coincides with the notion of Borel-fixed ideals. Ideals of the latter type are fixed by the action of the Borel subgroup of triangular matrices of the linear group and they correspond to the possible generic initial ideals by a famous result by Galligo [Gal74]. In the context of Hilbert schemes, Galligo's theorem states that each component and each intersection of components contains at least a point corresponding to a scheme defined by a Borel-fixed ideal. Hence, these ideals represent a promising tool for studying Hilbert schemes and to obtain a complete knowledge of their structure. To this aim, in recent years several authors [LR11, CR11, BCLR13, BLR13, LR13] developed algorithmic methods based on the use of strongly stable ideals to construct flat families corresponding to open subschemes of a Hilbert scheme. Once we are able to determine such open subschemes, in order to study all the components of a Hilbert scheme, we need to determine all its points corresponding to schemes defined by Borel-fixed ideals, i.e. all the saturated strongly stable ideals in a polynomial ring with a given Hilbert polynomial. The main feature of the package *StronglyStableIdeals* is a method to compute this set of ideals introduced in [CLMR11] and improved in [Lel12]. Several other tools are developed and presented in the paper.

1. Strongly stable ideals

Let $\mathbb{K}$ be any field and let $\mathbb{K}[\boldsymbol{x}] := \mathbb{K}[x_0, \dots, x_n]$ be the polynomial ring in $n+1$ variables with coefficients in $\mathbb{K}$. We assume $x_0 < x_1 < \cdots < x_n$. We use the multi-index notation to describe monomials, i.e. for every $\boldsymbol{\alpha} = (\alpha_0, \dots, \alpha_n) \in \mathbb{Z}_{\geqslant 0}^{n+1}$, $\boldsymbol{x}^{\boldsymbol{\alpha}} := x_0^{\alpha_0} \cdots x_n^{\alpha_n}$ and we denote by $\mathbb{T}_{n,s}$ the set of monomials of $\mathbb{K}[\boldsymbol{x}]$ of degree s. For any monomial $\boldsymbol{x}^{\boldsymbol{\alpha}}$, we denote by $\min \boldsymbol{x}^{\boldsymbol{\alpha}}$ and $\max \boldsymbol{x}^{\boldsymbol{\alpha}}$ the minimal and maximal index of a variable dividing $\boldsymbol{x}^{\boldsymbol{\alpha}}$.

Following [Gre10], we call *increasing* and *decreasing elementary moves* the elements of the field of fractions $\mathbb{K}(x_0, \dots, x_n)$

$$\mathrm{e}_i^+ := \frac{x_{i+1}}{x_i},\ i < n \qquad \text{and} \qquad \mathrm{e}_j^- := \frac{x_{j-1}}{x_j},\ j > 0.$$

We say that an elementary move $\mathrm{e}_i^{+/-}$ is *admissible* for a monomial $\boldsymbol{x}^{\boldsymbol{\alpha}}$ if $\mathrm{e}_i^{+/-}(\boldsymbol{x}^{\boldsymbol{\alpha}})$ is contained in $\mathbb{K}[\boldsymbol{x}]$, i.e. if $\alpha_i > 0$.

Definition 1.1. An ideal $I \subset \mathbb{K}[\boldsymbol{x}]$ is called *strongly stable* if

(1) I is a monomial ideal;
(2) for every $\boldsymbol{x}^{\boldsymbol{\alpha}} \in I$ and for every admissible increasing move e_i^+, the monomial $\mathrm{e}_i^+(\boldsymbol{x}^{\boldsymbol{\alpha}})$ in contained in I.

We recall that a strongly stable ideal is a Borel-fixed ideal. We now summarize some properties holding in general for Borel-fixed ideals and useful in this context.

Proposition 1.2 ([Gre10, Section 2])**.** *Let $I \subset \mathbb{K}[\boldsymbol{x}]$ be a strongly stable ideal.*

(i) The regularity of I is equal to the maximal degree of a generator (straightforward using the Eliahou-Kervaire resolution [EK90]).
(ii) Let $\mathfrak{m}$ be the irrelevant ideal of $\mathbb{K}[\boldsymbol{x}]$. Then, $(I : \mathfrak{m}) = (I : x_0)$, so that the ideal I is saturated if no generator involves the variable x_0.
(iii) The variable x_0 is a regular element for I, i.e. the multiplication by x_0 induces the short exact sequence

$$0 \longrightarrow \mathbb{K}[\boldsymbol{x}]/I(t-1) \xrightarrow{\cdot x_0} \mathbb{K}[\boldsymbol{x}]/I(t) \longrightarrow \mathbb{K}[\boldsymbol{x}]/(x_0, I)(t) \longrightarrow 0.$$

2010 *Mathematics Subject Classification.* 13P10, 13P99.
Key words and phrases. Strongly stable ideal, Borel-fixed ideal, Hilbert polynomial, Gotzmann number.

2. Hilbert polynomials

The Hilbert polynomial $p(t)$ of a homogeneous ideal $I \subset \mathbb{K}[\boldsymbol{x}]$ is the numerical polynomial such that for all s sufficiently large

$$\dim_{\mathbb{K}} \left(\mathbb{K}[\boldsymbol{x}]/I\right)_s = \dim_{\mathbb{K}} \left(\mathbb{K}[\boldsymbol{x}]_s/I_s\right) = \binom{n+s}{n} - \dim_{\mathbb{K}} I_s = p(s).$$

Obviously, not every numerical polynomial is a Hilbert polynomial of some homogenous ideal. Those being Hilbert polynomials have been completely described by Gotzmann [Got78].

Gotzmann's decomposition. *A numerical polynomial $p(t) \in \mathbb{Q}[t]$ is a Hilbert polynomial if, and only if, it can be decomposed as*

$$p(t) = \binom{n+a_1}{a_1} + \binom{n+a_2-1}{a_2} + \ldots + \binom{n+a_r-(r-1)}{a_r}, \qquad a_1 \geqslant a_2 \geqslant \cdots \geqslant a_r \geqslant 0. \tag{1}$$

This decomposition is strictly related to the Macaulay's decomposition

$$p(t) = \sum_{k=0}^{d} \left[\binom{t+k}{k+1} - \binom{t+k-m_k}{k+1} \right]$$

where $d = \deg p(t)$. For all $n \geqslant d+1$ the saturated lexicographic ideal in $\mathbb{K}[x_0, \ldots, x_n]$ with Hilbert polynomial $p(t)$ is

$$(x_n, \ldots, x_{d+2}, x_{d+1}^{b_d+1}, x_{d+1}^{b_d} x_d^{b_{d-1}+1}, \ldots, x_{d+1}^{b_d} x_d^{b_{d-1}} \cdots x_1^{b_0}),$$

where

$$b_d = \#\{a_j \mid a_j = d\} = m_d \quad \text{and} \quad b_k = \#\{a_j \mid a_j = k\} = m_k - m_{k+i},\ 0 \leqslant i < d.$$

The description of the lexicographic ideal in terms of the Gotzmann's decomposition suggests the following theorem.

Gotzmann's Regularity Theorem. *The regularity of a saturated ideal $I \subset \mathbb{K}[\boldsymbol{x}]$ with Hilbert polynomial $p(t)$ is at most r, where r is the number of terms in the decomposition (1) and it is called* Gotzmann number *of $p(t)$.*

Example 2.1. The package *StronglyStableIdeals* provides the method `isHilbertPolynomial` to determine if a numerical polynomial is a Hilbert polynomial.

```
Macaulay2, version 1.6
with packages: ConwayPolynomials, Elimination, IntegralClosure, LLLBases, PrimaryDecomposition,
               ReesAlgebra, TangentCone

i1 : loadPackage ("StronglyStableIdeals");

i2 : R = QQ[t]; -- the ring of numerical polynomials

i3 : p = 4*t;  q = 5*t-6;

i5 : isHilbertPolynomial p

o5 = true

i6 : isHilbertPolynomial q

o6 = false
```

Gotzmann's and Macaulay's decompositions of a Hilbert polynomial can be computed with the methods `gotzmannDecomposition` and `macaulayDecomposition`. These methods returns the list of the terms of the decomposition. The summand $\binom{t+e}{c}$ is constructed with the command `projectiveHilbertPolynomial(c,c-e)`.

```
i7 : gotzmannDecomposition p

o7 = {P , - P  + P , - 2*P  + P , - 3*P  + P , P , P }
       1     0    1       0    1       0    1   0   0

o7 : List

i8 : macaulayDecomposition p

o8 = {- P  + P , 7*P  - P , - P  + P , - 10*P  + 5*P  - P }
         0    1     0    1     1    2        0      1    2

o8 : List
```

Finally, the saturated lexicographic ideal L with Hilbert polynomial $p(t)$ in the polynomial ring $\mathbb{K}[\boldsymbol{x}]$ can be computed with the method `lexIdeal` and its regularity is equal to the Gotzmann number of $p(t)$.

```
i9 : L = lexIdeal(p,QQ[x,y,z,w])

             4 2   5
o9 = ideal (y z , y , x)

o9 : Ideal of QQ[x, y, z, w]

i10 : regularity L == gotzmannNumber p

o10 = true
```



FIGURE 1. The Borel sets defined in $\mathbb{T}_{2,3}$ and $\mathbb{T}_{2,4}$ by the ideal $(x_2^2, x_2x_1, x_1^4) \subset \mathbb{K}[x_0, x_1, x_2]$.

3. The main algorithm

In this section, we outline the strategy of the main algorithm. This algorithm was firstly described in [CLMR11] and then optimized in [Lel12]. The same problem has been previously discussed in [Ree92] and an alternative algorithm was later presented in [MN14].

We need to relate the properties of a strongly stable ideal with its Hilbert polynomial. If I is a strongly stable ideal, for each s the monomial basis of the homogeneous piece I_s of the ideal is a subset of $\mathbb{T}_{n,s}$ closed by increasing elementary moves. We call *Borel sets* such subsets of $\mathbb{T}_{n,s}$ (see Figure 1 for an example). Proposition 1.2*(i)* implies that for every saturated strongly stable ideal of $\mathbb{K}[\boldsymbol{x}]$ with Hilbert polynomial $p(t)$ and regularity at most s, the monomial basis of the homogeneous piece of degree s of the ideal is a Borel set with $p(s)$ elements, i.e. we can consider the following map

$$\left\{ \begin{array}{c} \text{saturated strongly stable ideals in } \mathbb{K}[\boldsymbol{x}] \text{ with} \\ \text{Hilbert polynomial } p(t) \text{ and regularity } \leqslant s \end{array} \right\} \hookrightarrow \left\{ \begin{array}{c} \text{Borel sets of } \mathbb{T}_{n,s} \\ \text{with } p(s) \text{ elements} \end{array} \right\}. \tag{2}$$

Moreover, Gotzmann's Regularity Theorem states that in order to consider all the saturated strongly stable ideals with a given Hilbert polynomial, it suffices to consider (2) for s equal to the Gotzmann number. Obviously, not every Borel set in $\mathbb{T}_{n,s}$ with $p(s)$ elements corresponds to an ideal with Hilbert polynomial $p(t)$. To identify the image of the previous map, we recall a definition and a proposition by Mall.

Definition 3.1 ([Mal97, Definition 2.7])**.** Let $B \subset \mathbb{T}_{n,s}$ be a Borel set. The set $B^{(i)} := \{\boldsymbol{x}^{\boldsymbol{\alpha}} \in B \mid \min \boldsymbol{x}^{\boldsymbol{\alpha}} = i\}$ is called *i-growth class* of B. The sequence $\mathrm{gv}(B) := (|B^{(0)}|, \dots, |B^{(n)}|)$ is called *growth vector* of B.

Proposition 3.2 ([Mal97, Proposition 3.2])**.** *Let $I \subset \mathbb{K}[\boldsymbol{x}]$ be a strongly stable ideal generated by the monomials of a Borel set $B \subset \mathbb{T}_{n,s}$ and let $p(t)$ be its Hilbert polynomial. Then,*

$$p(t) = \binom{n+t}{n} - \sum_{k=0}^{n} \left|B^{(k)}\right| \binom{k+t-s}{k}, \qquad \forall\, t \geqslant s. \tag{3}$$

Now, the point is to determine the growth vector of a Borel set $B \subset \mathbb{T}_{n,s}$ starting from the Hilbert polynomial. The first difference polynomial is

$$\begin{aligned} \Delta p(t) = p(t) - p(t-1) &= \binom{n+t}{n} - \binom{n+t-1}{n} - \sum_{k=0}^{n} \left|B^{(k)}\right| \left[\binom{k+t-s}{k} - \binom{k+t-s-1}{k}\right] = \\ &= \binom{n+t-1}{n-1} - \sum_{k=1}^{n} \left|B^{(k)}\right| \binom{k+t-s-1}{k-1} \end{aligned}$$

so that repeating this computation and evaluating these polynomials for $t = s$, we have

$$\sum_{k=i}^{n} \left|B^{(k)}\right| = \binom{n+s-i}{n-i} - \Delta^i p(s), \qquad \forall\, i = 0, \dots, n,$$

where $\Delta^0 p(t) = p(t)$ and $\Delta^i p(t) = \Delta^{i-1} p(t) - \Delta^{i-1} p(t-1)$ (notice that $\Delta^i p(t) = 0$ for $i > \deg p(t)$). These conditions allow to deduce that $|B^{(n)}| = 1$ and for $i < n$

$$\left|B^{(i)}\right| = \sum_{k=i}^{n} \left|B^{(k)}\right| - \sum_{k=i+1}^{n} \left|B^{(k)}\right| = \binom{n+s-i-1}{n-i} - \Delta^i p(s) + \Delta^{i+1} p(s).$$

Let us call the sequence of such integers *growth vector* of $p(t)$ in degree s and let us denote it by $\mathrm{gv}_s\big(p(t)\big)$.



FIGURE 2. On the left the Borel set corresponding to the ideal $(x_2^3, x_2^2x_1, x_2x_1^4)$ and on the right the one corresponding to $(x_2^3, x_2^2x_1^2, x_2x_1^3)$. The dashed lines separate the growth classes of the Borel sets.

Proposition 3.3 (cf. [Lel12, Theorem 3.3]). *Let $p(t)$ be a Hilbert polynomial. There is a bijective map*

$$\text{(4)}\qquad \begin{array}{ccc} \left\{\begin{array}{c}\textit{saturated strongly stable ideals}\\ \textit{in } \mathbb{K}[\boldsymbol{x}] \textit{ with Hilbert polynomial}\\ p(t) \textit{ and regularity } \leqslant s\end{array}\right\} & \overset{1:1}{\longleftrightarrow} & \left\{\begin{array}{c}\textit{Borel sets of } \mathbb{T}_{n,s}\\ \textit{with } p(s) \textit{ elements and}\\ \textit{growth vector } \mathrm{gv}_s\big(p(t)\big)\end{array}\right\} \\ I & \longrightarrow & \textit{monomial basis of } I_s \\ \textit{saturation of } (B) & \longleftarrow & B \end{array}$$

In order to determine the Borel sets of Proposition 3.3, we use a recursive algorithm based on Proposition 1.2*(iii)*. Indeed, if $I \subset \mathbb{K}[x_0, \dots, x_n]$ is a strongly stable ideal with Hilbert polynomial $p(t)$ and B is the associated Borel set in $\mathbb{T}_{n,s}$, then the subset $B' = \{\boldsymbol{x}^{\boldsymbol{\alpha}} \in B \mid \min \boldsymbol{x}^{\boldsymbol{\alpha}} > 0\} \subset B$ is a Borel set in $\mathbb{T}_{n-1,s}$ corresponding to the strongly stable ideal $I' = (x_0, I) \cap \mathbb{K}[x_1, \dots, x_n] \subset \mathbb{K}[x_1, \dots, x_n]$ with Hilbert polynomial $\Delta p(t)$.

Example 3.4. We want to find the strongly stable ideals of the polynomial ring $\mathbb{K}[x_0, x_1, x_2]$ with regularity at most 5 defining schemes with Hilbert polynomial $p(t) = t+6$. The Gotzmann number of $p(t)$ is 6 and its growth vector in degree 5 is $\mathrm{gv}_5(t+6) = (5, 4, 1)$. In order to accomplish our task, we start considering the strongly stable ideals of $\mathbb{K}[x_1, x_2]$ with Hilbert polynomial $\Delta p(t) = 1$ and regularity at most 5 corresponding to Borel sets with growth vector $\mathrm{gv}_5(\Delta p(t)) = (4, 1)$. There is a unique Borel set $B' = \{x_2^5, x_2^4x_1, x_2^3x_1^2, x_2^2x_1^3, x_2x_1^4\}$. Since x_1^5 is not contained in B', any Borel set $B \subset \mathbb{T}_{2,5}$ with growth vector $(5, 4, 1)$ could not contain the monomials that can be obtained from x_1^5 by applying decreasing elementary moves, i.e. $x_1^4x_0$, $x_1^3x_0^2$, $x_1^2x_0^3$, $x_1x_0^4$ and x_0^5. In Figure 2, there are the Borel sets and the corresponding strongly stable ideals with Hilbert polynomial $t+6$.

This package provides the method `stronglyStableIdeals` to compute the set of strongly stable ideals of a given polynomial ring with fixed Hilbert polynomial and bounded regularity.

```
i11 : stronglyStableIdeals(4*t,QQ[x,y,z,w])

                    5   4 2                    2   4    5                2   2  4                2   3
o11 = {ideal (x, y , y z ), ideal (x*z, x*y, x , y z, y ), ideal (x*y, x , x*z , y ), ideal (x*y, x , y )}

o11 : List

i12 : stronglyStableIdeals(4*t,QQ[x,y,z,w],MaxRegularity=>4)

                    2   2  4                2   3
o12 = {ideal (x*y, x , x*z , y ), ideal (x*y, x , y )}

o12 : List
```

4. Segment ideals

The transitive closure of the order relation

$$\text{(5)}\qquad \boldsymbol{x}^{\boldsymbol{\alpha}} >_B \boldsymbol{x}^{\boldsymbol{\beta}} \iff \boldsymbol{x}^{\boldsymbol{\beta}} = \mathrm{e}_i^-(\boldsymbol{x}^{\boldsymbol{\alpha}})$$

induces a partial order on the set of monomials of any degree called *Borel order*. Every graded term ordering turns out to be a refinement of this partial order. Since a Borel set B is closed with respect to the Borel order, i.e. $\boldsymbol{x}^{\boldsymbol{\alpha}} >_B \boldsymbol{x}^{\boldsymbol{\beta}}$, $\boldsymbol{x}^{\boldsymbol{\beta}} \in B \Rightarrow \boldsymbol{x}^{\boldsymbol{\alpha}} \in B$, it is natural to ask whether there exists a term ordering $\prec$ preserving the same property. For instance, for the lexicographic ideal, the graded lexicographic order separates, in each degree, the monomials contained in the ideal from those not contained in it. In [CLMR11] several notions of segment ideals are introduced.

Definition 4.1 ([CLMR11, Definitions 3.1, 3.7]). A Borel set $B \subset \mathbb{T}_{n,s}$ is called *segment* if there exists a term ordering $\prec$ such that $\boldsymbol{x}^{\boldsymbol{\alpha}} \succ \boldsymbol{x}^{\boldsymbol{\beta}}$, for all $\boldsymbol{x}^{\boldsymbol{\alpha}} \in B$ and $\boldsymbol{x}^{\boldsymbol{\beta}} \in \mathbb{T}_{n,s} \setminus B$.

Let $I \subset \mathbb{K}[\boldsymbol{x}]$ be a saturated strongly stable ideal.

(i) I is called *hilb-segment* if the Borel set $I \cap \mathbb{T}_{n,r}$ is a segment, where r is the Gotzmann number of the Hilbert polynomial of I.
(ii) I is called *reg-segment* if the Borel set $I \cap \mathbb{T}_{n,m}$ is a segment, where m is the regularity of I.
(iii) I is called *gen-segment* if there exists a term ordering $\prec$ such that $\boldsymbol{x}^{\alpha} \succ \boldsymbol{x}^{\boldsymbol{\beta}}$ for each minimal generator $\boldsymbol{x}^{\boldsymbol{\alpha}}$ of degree s of I and for all $\boldsymbol{x}^{\boldsymbol{\beta}} \in \mathbb{T}_{n,s} \setminus I_s$.

These notions are very important in the construction of flat families based on properties of Gröbner bases and in general for the study of the Hilbert scheme. The package provides three methods for determining whether a strongly stable ideal may be some type of segment (and, in case, gives the term ordering). This methods use tools of the package *gfanInterface* so that the term ordering is given as the corresponding weight vector of the variables.

```
i13 : SSI = stronglyStableIdeals(7,3,MaxRegularity=>5)

                2     2   5          2   4     3          2   2     3   4
o13 = {ideal (x , x x , x ), ideal (x , x , x x ), ideal (x x , x x , x , x )}
                2   2 1   1          2   1   2 1          2 1   2 1   2   1

o13 : List

i14 : isHilbSegment(SSI#0)

o14 = (true, {7, 3, 1})

o14 : Sequence

i15 : isRegSegment(SSI#1)

o15 = (false, )

o15 : Sequence

i16 : isGenSegment(SSI#1)

o16 = (true, {4, 3, 1})

o16 : Sequence
```

References

[BCLR13] Cristina Bertone, Francesca Cioffi, Paolo Lella and Margherita Roggero, *Upgraded methods for the effective computation of marked schemes on a strongly stable ideal*, J. Symbolic Comput. **50** (2013), 263–290.

[BLR13] Cristina Bertone, Paolo Lella and Margherita Roggero, *A Borel open cover of the Hilbert scheme*, J. Symbolic Comput. **53** (2013), 119–135.

[CLMR11] Francesca Cioffi, Paolo Lella, Maria Grazia Marinari and Margherita Roggero, *Segments and Hilbert schemes of points*, Discrete Math. **311** (2011), no. 20, 2238–2252.

[CR11] Francesca Cioffi and Margherita Roggero, *Flat families by strongly stable ideals and a generalization of Gröbner bases*, J. Symbolic Comput. **46** (2011), no. 9, 1070–1084.

[EK90] Shalom Eliahou and Michel Kervaire, *Minimal resolutions of some monomial ideals*, J. Algebra **129** (1990), no. 1, 1–25.

[Gal74] André Galligo, *À propos du théorème de-préparation de Weierstrass*, Fonctions de plusieurs variables complexes (Sém. François Norguet, octobre 1970–décembre 1973; à la mémoire d'André Martineau), Lecture Notes in Math., vol. 409, Springer, Berlin, 1974, Thèse de 3ème cycle soutenue le 16 mai 1973 à l'Institut de Mathématique et Sciences Physiques de l'Université de Nice, pp. 543–579.

[Got78] Gerd Gotzmann, *Eine Bedingung für die Flachheit und das Hilbertpolynom eines graduierten Ringes*, Math. Z. **158** (1978), no. 1, 61–70.

[Gre10] Mark L. Green, *Generic initial ideals*, Six lectures on commutative algebra, Mod. Birkhäuser Class., Birkhäuser Verlag, Basel, 2010, pp. 119–186.

[Lel12] Paolo Lella, *An efficient implementation of the algorithm computing the Borel-fixed points of a Hilbert scheme*, ISSAC 2012 — Proceedings of the 37th International Symposium on Symbolic and Algebraic Computation (2012), 242–248.

[LR11] Paolo Lella and Margherita Roggero, *Rational components of Hilbert schemes*, Rend. Semin. Mat. Univ. Padova **126** (2011), 11–45.

[LR13] ———, *On the functoriality of marked families*, ArXiv e-prints (arxiv.org/abs/1307.7657) (2013).

[Mal97] Daniel Mall, *Betti numbers, Castelnuovo Mumford regularity, and generalisations of Macaulay's theorem*, Comm. Algebra **25** (1997), no. 12, 3841–3852.

[Mal00] ———, *Connectedness of Hilbert function strata and other connectedness results*, J. Pure Appl. Algebra **150** (2000), no. 2, 175–205.

[MN14] Dennis Moore and Uwe Nagel, *Algorithms for strongly stable ideals*, Math. Comp. (2014).

[Ree92] Alyson A. Reeves, *Combinatorial structure on the Hilbert scheme*, Ph.D. thesis, Cornell University, 1992.

Davide Alberelli, University of Osnabrück, Institut für Mathematik, Albrechtstr. 28a, 49076 Osnabrück, Germany.

E-mail address: dalberelli@uos.de

URL: http://www.mathematik.uni-osnabrueck.de/graduiertenkolleg/personen/alberelli_davide.html

Paolo Lella, Dipartimento di Matematica dell'Università degli Studi di Torino, Via Carlo Alberto 10, 10123 Torino, Italy.

E-mail address: paolo.lella@unito.it

URL: http://www.personalweb.unito.it/paolo.lella/